

**CURRENT CONCEPTS REVIEW**

# Radiofrequency Ablation for the Management of Pain of Spinal Origin in Orthopedics

E. Carlos Rodriguez-Merchan, MD, PhD; Alberto D. Delgado-Martinez, MD, PhD;  
Javier De Andres-Ares, MD, PhD

Research performed at Department of Orthopedic Surgery, La Paz University Hospital-IdiPaz, Madrid, Spain

Received: 21 March 2023

Accepted: 28 August 2023

**Abstract**

Radiofrequency (RF) is a minimally invasive procedure for interrupting or modifying nociceptive pathways to manage musculoskeletal neuropathic and nociplastic pain. RF has been used to manage cervical and lumbar facet joint syndromes. The benefits of RF are the following: it is safer than surgery; it doesn't need general anesthesia, by that means diminishing the complications; it causes pain alleviation for a minimum of 3-4 months; it can be repeated if needed; it improves joint function; and it minimizes the necessity for oral pain treatment. RF must not be used in pregnant women; unstable joints, uncontrolled diabetes mellitus; and presence of an implanted defibrillator. Even though complications from RF are rare, possible adverse events are infection, hemorrhage, numbness or dysesthesia, greater pain in the procedure area, and differentiation impact. Even though there is a peril of injuring non-targeted neural tissue and other tissues, this can be lessened by carrying out the procedure with the help of fluoroscopy, ultrasonography or computed tomography. RF seems to be a helpful procedure for relieving chronic pain syndromes; despite this, definite proof of the procedure's effectiveness is still needed. RF is an encouraging procedure for treating chronic spinal pain, especially when other procedures are pointless or not feasible.

**Level of evidence:** III

**Keywords:** Chronic pain, Management, Radiofrequency, Spine

**Introduction**

**R**adiofrequency (RF) waves destroy the neural tissue that sends pain impulses to the brain. An RF circuit includes an RF generator creating the energy, a needle that is introduced into the individual, and a ground pad that is connected to dispel the energy as long as it is attached to the RF generator. RF energy is applied and sent to the individual, which makes the ions in the transmitting tissues to swing, liberating heat energy at the tip of the needle but without making the needle hot. The oscillations of the ions make the encircling tissue of the tip of the needle hot.<sup>1,2</sup>

The utilization of RF (also known as rhizotomy or neurotomy) for treating chronic pain was first published in 1931.<sup>3</sup> In 1950s, the first commercial RF generator was created.<sup>4</sup> Publications of other RF methods appeared in 1998, when pulsed RF was developed to cause a less "aggressive" technique.<sup>5</sup> The way by which pulsed RF

alleviates pain endures unsure. A third category of RF ("cooled" RF) started utilization in the mid-1990s. It puts cold saline solution to the tip of the RF probe, which produces extensive, controlled injuries;<sup>6-10</sup> nonetheless, it was not until 2008 that the first publications mentioning the utilization of cooled RF were reported.<sup>11-14</sup> Since then, cooled RF has been demonstrated to be valuable for many indications.<sup>15-24</sup>

The objective of this paper is to perform a narrative review of current publications on the capacity of RF in managing spinal pain. For this aim, a quest was carried in PubMed on July 15, 2023, utilizing "radiofrequency facet joint" as key words. We encountered 464 publications, 46 of which were chosen because, in our belief, they were the most relevant and were absolutely connected to the title of this paper. Therefore, 418 were excluded. Besides, 3 chapters from a relevant book published on the topic were analyzed,

**Corresponding Author:** E. Carlos Rodriguez-Merchan, Department of Orthopedic Surgery, La Paz University Hospital-IdiPaz, Madrid, Spain

Email: ecrmerchan@hotmail.com



THE ONLINE VERSION OF THIS ARTICLE  
ABJS.MUMS.AC.IR

amounting to a total of 49 references. Consequently, this paper is not a systematic review of the literature but rather a narrative review.


### **Indications, contraindications and advantages of RF**

RF is advised for treating pain emerging from various musculoskeletal origins only if conservative management [pain killers, nonsteroidal anti-inflammatory drugs (NSAIDs), weight loss, Physical and Rehabilitation Medicine] has failed to render pain relief and the individual is not a candidate for surgical treatment.<sup>25-27</sup> RF must not be used in pregnant women, individuals with uncontrolled diabetes mellitus, and in those who wear an internal defibrillator. RF is safer than surgery, does not need general anaesthesia (therefore reducing complications), renders pain alleviation for a minimum of 3-4 months, can be done again if required; ameliorates joint function, and minimizes the necessity of oral pain treatment.<sup>15-24</sup>

### **RF in cervical facet joint syndrome**

History, physical exam, and radiographic evaluation might help detect an individual's candidacy for diagnostic cervical medial branch nerve blocks as a reasonable step in the confirmation of cervical facet joint pain, as well as to evaluate their candidacy for subsequent RF. The confirmation of cervical facet-mediated pain typically should be made by means of local anaesthetic blocks, either an intraarticular injection of the facet joint or a block of the nerves innervating the facet joint (i.e., the dorsal primary rami or medial branches). Before performing RF of the cervical facet, we should perform "double blocks" to confirm that the cervical facet joint is the primary pain generator.<sup>28-35</sup>

In a systematic review, successful pain alleviation ranged from 30% to 50%.<sup>36</sup> [Figure 1] shows fluoroscopically guided conventional RF for cervical medial branch denervation.



**Figure 1.** Fluoroscopically guided conventional radiofrequency (RF) for cervical medial branch denervation: (A) anteroposterior fluoroscopic view; (B) lateral fluoroscopic image

### **RF in lumbar zygapophysial (facet) joint syndrome**

Lumbar facet joints are the most frequent cause of chronic low back pain (compared with discs and sacroiliac joints), with an incidence of 21%-41%, and are to a greater degree the most frequent cause of chronic low back pain in individuals older than 65 years. The diagnosis of lumbar facet joint syndrome in individuals with low back pain must be

confirmed by means of diagnostics blocks.<sup>37-41</sup>

Afifi et al compared the RF of lumbar facet joints with chemical neurolysis using ethyl alcohol 95% (EA-95) or glycerol 20% (Gly-20).<sup>42</sup> Ninety-five individuals with a mean age of 68 years were analyzed, of whom 30 experienced RF, 30 were managed with EA-95, and 35 were managed with Gly-20. After 6 weeks, the RF individuals had substantially lower visual analogue scale (VAS) scores compared with the EA-95 cohort. After 6 months, the VAS was substantially inferior in the RF individuals than in the Gly-20 cohort. One year after the procedure, the VAS scores were substantially inferior in the RF cohort compared with the Gly-20 cohort. This study demonstrated that RF was associated with improved pain alleviation and quality of life (QoL) compared with chemical neurolysis. RF should therefore be deemed as the management of choice for individuals with chronic low back pain due to lumbar facet joint syndrome.<sup>42</sup>

Li et al published a systematic review to assess and compare the efficacy of various RF methods for treating facet joint-derived chronic low back pain.<sup>43</sup> Procedures included typical RF denervation, pulsed RF (PRF) denervation, PRF treatment of the dorsal root ganglia (PRF-DRG), RF facet capsule denervation (RF-FC), and RF ablation under endoscopic guidance (ERFA). The outcome was the modification in VAS average score from baseline at  $\leq 6$  months and 12 months. Moderate evidence showed that conventional RF denervation had a bigger impact on pain alleviation than the sham control in the short-run and long term. Fair evidence demonstrated that PRF denervation was more efficacious than the sham control for pain over the long run. ERFA denervation was more efficacious for pain alleviation than the sham control in the short run and the long term. Fair evidence demonstrated that RF-FC denervation was more efficacious for pain alleviation than the sham control in the long term. PRF-DRG denervation was more efficacious for pain alleviation than the sham control in the short run. RF was an efficacious alternative for individuals with facet joint-derived chronic low back pain.<sup>41</sup> Turtle et al reported that instrumented lumbar spinal fusion can produce facet-mediated pain at levels adjacent to the surgical site due to disruption of lumbar medial branch nerve(s) and that RF could be useful in these cases.<sup>44</sup>

Even though complications from cervical and lumbar RF are rare, possible adverse events can include infection, hemorrhage, numbness or dysesthesias, greater pain in the area of the procedure, and deafferentation effects. Dysesthesia (a "sunburn"-type sensation) can occur after cervical RF, which has been encountered to be self-solving. Even though the peril for cervical nerve root injury exists, it can be lessened by performing the procedure with the help of fluoroscopy, and sensory and motor testing prior to lesioning.<sup>45</sup> displays fluoroscopically guided conventional RF for lumbar medial branch denervation.

According to Lo Bianco et al, lumbar facet joint syndrome is one of the main causes of chronic low back pain. RF ablation utilizing a V-shaped active tip needle might produce a greater injury in the medial branch nerves, ameliorating clinical result. They assessed the effectiveness and the possibility of

RF ablation utilizing V-shaped active tip needles. The inclusion criteria were the following: adult individuals (>18 years), diagnosis of chronic lumbar facet joint pain and unsuccessful attempt of conservative management. The exclusion criteria were the following: lumbar pain unrelated to facet joints and prior spinal/lumbar surgical procedures. The primary outcome of the analysis was a modification in pain intensity at follow-up. The secondary parameters were QoL, the occurrence of complications and the influence on postprocedural analgesic consumption. For these objectives, the following parameters were evaluated: pre- and posttreatment numeric rating scale (NRS), neuropathic pain 4 questions (DN4), EuroQoL - EQ-5D-3L, EQ-VAS, EQ-index and North American Spine Society (NASS) index. Sixty-four individuals were included. 7.8% of individuals at 1-month, 37.5% at 3-month, 40.6% at 6-month and 35.9% at 9-month follow-up reported a decrease of more than 80% in NRS, DN4, EQ-index and EQ-5D-VAS at the different time-points. This study demonstrated that RF ablation utilizing a V-shaped active tip needle may be a possible and efficacious management for chronic lumbar facet joint syndrome.<sup>46</sup>




Figure 2. Fluoroscopically guided conventional radiofrequency (RF) for lumbar medial branch denervation: (a) anteroposterior fluoroscopic view; (b) lateral fluoroscopic image

#### RF in sacroiliac joint pain

In a systematic review, Maccagnano et al compared conventional RF and cooled RF to treat sacroiliac joint pain. Nine studies were identified, with 276 individuals suffering from sacroiliac joint pain that were managed with RF. The two techniques showed no major adverse events, and no statistical difference between the two procedures studied was found. The authors stated that bibliography on the subject is currently missing, and well-constructed randomized clinical trials (RCTs) are required to assess this deficiency.<sup>47</sup> shows fluoroscopically guided conventional RF for sacroiliac joint denervation.

Maalouly and Rao have recently analyzed 81 individuals (59 women, 22 men) who experienced cooled RF ablation. Their average age was 55 years. Twenty-two of the individuals had experienced fusions of the lumbar spine in previous years. Follow-up period ranged from 6 months to 1.5 years. After RF ablation, seven individuals progressed to fusions, and 6 individuals had to have the surgical technique repeated to alleviate their pain. This study concluded that sacroiliac joint RF ablation was a good alternative in the management of sacroiliac joint pain demonstrating good outcomes in the

short-run follow-up period. It is an easy technique that can be performed in less than 30 minutes and is able of rendering substantial pain alleviation for individuals with sacroiliac joint pain.<sup>48</sup>




Figure 3. Fluoroscopically guided conventional radiofrequency (RF) for sacroiliac joint denervation: (A) clinical view; (B) anteroposterior fluoroscopic view; (C) lateral fluoroscopic image; (D) oblique fluoroscopic view

#### RF in coccydynia

Through a 2022 systematic review, Andersen et al evaluated the effectiveness of accessible management alternatives for patients with chronic coccydynia.<sup>45</sup> The greatest reduction in pain was experienced by the individuals who experienced RF treatment. An equivalent amelioration was experienced with extracorporeal shockwave therapy, coccygectomy, and injection. Although pain improved in all the studied alternatives, the average pain improvement was less for those who underwent ganglion block, stretching/manipulation and conservative/usual care. This report emphasized that treatment of coccydynia must begin with non-invasive approaches prior to considering coccygectomy. Nonsurgical treatment provided pain relief for many of the patients. Coccygectomy might be helpful for refractory patients.<sup>49</sup>

#### Conclusion

RF is a minimally invasive procedure utilized to destroy or modify nociceptive pathways for managing spinal pain. Satisfactory outcomes have been observed in sacroiliac joint pain utilizing RF of the lateral branches of the L4-S3 dorsal rami. In cervical and lumbar facet joint syndromes, RF of the respective branch nerves has yielded promising results. RF seems to be a helpful procedure for relieving chronic spinal syndromes, even if, until now, solid proof is still needed on

the effectiveness of this procedure. RF is a hopeful procedure for treating chronic spinal pain, especially when other procedures are ineffective or not feasible.

### Acknowledgement

Not applicable

**Conflict of interest:** None

**Funding:** None

E. Carlos Rodriguez-Merchan MD, PhD <sup>1</sup>

Alberto D. Delgado-Martinez PhD <sup>2,3</sup>  
Javier De Andres-Ares MD <sup>4</sup>

1 Department of Orthopedic Surgery, La Paz University Hospital-IdiPaz, Madrid, Spain

2 Department of Orthopedic Surgery, Hospital Universitario de Jaén, Jaén, Spain

3 Department of Surgery, University of Jaén, Jaén, Spain

4 Pain Unit, Department of Anesthetics, La Paz University Hospital, Madrid, Spain

### References

- Choi EJ, Choi YM, Jang EJ, Kim JY, Kim TK, Kim KH. Neural ablation and regeneration in pain practice. *Korean J Pain*. 2016; 29(1):3-11. doi: 10.3344/kjp.2016.29.1.3.
- Rodriguez-Merchan EC, Delgado Martinez AD, De Andres-Ares J. Upper limb and lower limb radiofrequency treatments in orthopaedics. *EFORT Open Rev* 2023; 8(6):424-435. doi: 10.1530/EOR-22-0127.
- Hagedorn JM, Golovac S, Deer TR, Azeem N. History and development of radiofrequency ablation for chronic pain. In: *Essentials of Radiofrequency Ablation of the Spine and Joints*. 1th ed. Deer TR, Azeem N, eds. Springer Nature Switzerland AG, Cham; 2021.
- Aronow S. The use of radiofrequency power in making lesions in the brain. *J Neurosurg*. 1960; 17:431-438. doi: 10.3171/jns.1960.17.3.0431.
- van Suijlekom HA, Weber WE, van Kleef M, Barendse GA, Sluijter ME. Radiofrequency cervical zygapophyseal joint neurotomy for cervicogenic headache: a short term follow-up study. *Funct Neurol*. 1998; 13(1):82-83.
- Nakagawa H, Yamanashi WS, Pitha JV, et al. Comparison of in vivo tissue temperature profile and lesion geometry for radiofrequency ablation with a saline-irrigated electrode versus temperature control in a canine thigh muscle preparation. *Circulation*. 1995; 91(8):2264-2273. doi: 10.1161/01.cir.91.8.2264.
- Ruffy R, Imran MA, Santel DJ, Wharton JM. Radiofrequency delivery through a cooled catheter tip allows the creation of larger endomyocardial lesions in the ovine heart. *J Cardiovasc Electrophysiol*. 1995; 6(12):1089-1096. doi: 10.1111/j.1540-8167.1995.tb00386.x.
- Buscarini L, Rossi S. Technology for radiofrequency thermal ablation of liver tumors. *Semin Laparosc Surg*. 1997; 4(2):96-101. doi: 10.1053/SLAS00400096.
- Solbiati L, Goldberg SN, Ierace T, et al. Hepatic metastases: percutaneous radio-frequency ablation with cooled-tip electrodes. *Radiology*. 1997; 205(2):367-373. doi: 10.1148/radiology.205.2.9356616.
- Petersen HH, Chen X, Pietersen A, Svendsen JH, Haunsø S. Lesion size in relation to ablation site during radiofrequency ablation. *Pacing Clin Electrophysiol*. 1998; 21(1 Pt 2):322-326. doi: 10.1111/j.1540-8159.1998.tb01114.x.
- Cohen SP, Hurley RW, Buckenmaier CC, Kurihara C, Morlando B, Dragovich A. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology*. 2008; 109(2):279-288. doi: 10.1097/ALN.0b013e31817f4c7c.
- Kapural L, Nageeb F, Kapural M, Cata JP, Narouze S, Mekhail N. Cooled radiofrequency system for the treatment of chronic pain from sacroiliitis: the first case-series. *Pain Pract*. 2008; 8(5):348-354. doi: 10.1111/j.1533-2500.2008.00231.x.
- Kapural L, Ng A, Dalton J, et al. Intervertebral disc biacuplasty for the treatment of lumbar discogenic pain: results of a six-month follow-up. *Pain Med*. 2008; 9(1):60-67. doi: 10.1111/j.1526-4637.2007.00407.x.
- Kapural L. Intervertebral disk cooled bipolar radiofrequency (intradiskal biacuplasty) for the treatment of lumbar diskogenic pain: a 12-month follow-up of the pilot study. *Pain Med*. 2008; 9(4):407-408. doi: 10.1111/j.1526-4637.2008.00464.x.
- Vu T, Chhatre A. Cooled radiofrequency ablation for bilateral greater occipital neuralgia. *Case Rep Neurol Med*. 2014; 2014:257373. doi: 10.1155/2014/257373.
- McCormick ZL, Walker J, Marshall B, McCarthy R, Walega DR. A novel modality for facet joint denervation: cooled radiofrequency ablation for lumbar facet syndrome. A case series. *Phys Med Rehabil Int*. 2014; 1(5):5.
- Walega D, Roussis C. Third-degree burn from cooled radiofrequency ablation of medial branch nerves for treatment of thoracic facet syndrome. *Pain Pract*. 2014; 14(6):e154-e158. doi: 10.1111/papr.12222.
- Bellini M, Barbieri M. Cooled radiofrequency system relieves chronic knee osteoarthritis pain: the first case-series. *Anaesthesiol Intensive Ther*. 2015; 47(1):30-33. doi: 10.5603/AIT.2015.0003.
- Menzies RD, Hawkins JK. Analgesia and improved performance in a patient treated by cooled radiofrequency for pain and dysfunction postbilateral total knee replacement. *Pain Pract*. 2015; 15(6):e54-e58. doi: 10.1111/papr.12292.
- Bajaj PS, Napolitano J, Wang W, Cheng J, Singh JR. Cooled versus conventional thermal radiofrequency neurotomy for the treatment of lumbar facet-mediated pain. *PM R*. 2015; 7(10):1095-1101. doi: 10.1016/j.pmrj.2015.09.001.
- Reddy RD, McCormick ZL, Marshall B, Mattie R, Walega DR. Cooled radiofrequency ablation of genicular nerves for knee osteoarthritis pain: a protocol for patient selection and case series. *Anesth Pain Med*. 2016; 6(6):e39696. doi: 10.5812/aapm.39696.

22. Kapural L, Jolly S, Mantoan J, Badhey H, Ptacek T. Cooled radiofrequency Neurotomy of the articular sensory branches of the obturator and femoral nerves - combined approach using fluoroscopy and ultrasound guidance: technical report, and observational study on safety and efficacy. *Pain Physician*. 2018; 21(3):279-284.
23. McCormick ZL, Choi H, Reddy R, et al. Randomized prospective trial of cooled versus traditional radiofrequency ablation of the medial branch nerves for the treatment of lumbar facet joint pain. *Reg Anesth Pain Med*. 2019; 44(3):389-397. doi: 10.1136/rapm-2018-000035.
24. Naber J, Lee N, Kapural L. Clinical efficacy assessment of cooled radiofrequency ablation of the hip in patients with avascular necrosis. *Pain Manag*. 2019; 9(4):355-359. doi: 10.2217/pmt-2018-0083.
25. Azadinia F, Ebrahimi-Takamjani I, Kamyab M, Asgari M, Parnianpour M. Immediate effects of lumbosacral orthosis on postural stability in patients with low back pain: a preliminary study. *Arch Bone Jt Surg*. 2019; 7(4):360-366.
26. Pourahmadi M, Sahebalam M, Bagheri R. Effectiveness of proprioceptive neuromuscular facilitation on pain intensity and functional disability in patients with low back pain: a systematic review and meta-analysis. *Arch Bone Jt Surg*. 2020; 8(4):479-501. doi: 10.22038/abjs.2020.45455.2245.
27. Zamani H, Dadgou M, Akbari M, Sarrafzadeh J, Pourahmadi M. Effects of external focus and motor control training in comparison with motor control training alone on pain, thickness of trunk muscles and function of patients with recurrent low back pain: a single blinded, randomized controlled trial. *Arch Bone Jt Surg*. 2022; 10(9):766-774. doi: 10.22038/ABJS.2022.56938.2824.
28. Ghosh P, Shah J, Esposito M, Azeem N, Deer TR. Cervical Spine. In: *Essentials of Radiofrequency Ablation of the Spine and Joints*. 1st ed. Deer TR, Azeem N, eds. Springer Nature Switzerland AG, Cham; 2021.
29. Holz SC, Sehgal N. What is the correlation between facet joint radiofrequency outcome and response to comparative medial branch blocks? *Pain Physician*. 2016; 19(3):163-172.
30. Cohen SP, Strassels SA, Kurihara C, et al. establishing an optimal "cutoff" threshold for diagnostic lumbar facet blocks: a prospective correlational study. *Clin J Pain*. 2013; 29(5):382-391. doi: 10.1097/AJP.0b013e31825f53bf.
31. Derby R, Melnik I, Lee JE, Lee SH. Correlation of lumbar medial branch neurotomy results with diagnostic medial branch block cutoff values to optimize therapeutic outcome. *Pain Med*. 2012; 13(12):1533-1546. doi: 10.1111/j.1526-4637.2012.01500.x.
32. Cohen SP, Williams KA, Kurihara C, et al. Multicenter, randomized, comparative cost-effectiveness study comparing 0, 1, and 2 diagnostic medial branch (facet joint nerve) block treatment paradigms before lumbar facet radiofrequency denervation. *Anesthesiology*. 2010; 113(2):395-405. doi: 10.1097/ALN.0b013e3181e33ae5.
33. Kweon TD, Kim JY, Lee HY, Kim MH, Lee YW. Anatomical analysis of medial branches of dorsal rami of cervical nerves for radiofrequency thermocoagulation. *Reg Anesth Pain Med*. 2014; 39(6):465-471. doi: 10.1097/AAP.0000000000000175.
34. Manchikanti L, Hirsch JA, Kaye AD, Boswell MV. Cervical zygapophysial (facet) joint pain: effectiveness of interventional management strategies. *Postgrad Med*. 2016; 128(1):54-68. doi: 10.1080/00325481.2016.1105092.
35. Manchikanti L, Kaye AD, Soin A, et al. Comprehensive evidence-based guidelines for facet joint interventions in the management of chronic spinal pain: American Society of Interventional Pain Physicians (ASIPP) guidelines facet joint interventions 2020 guidelines. *Pain Physician*. 2020; 23(3S):S1-S127.
36. Suer M, Wahezi SE, Abd-Elsayed A, Sehgal N. Cervical facet joint pain and cervicogenic headache treated with radiofrequency ablation: a systematic review. *Pain Physician*. 2022; 25(3):251-263.
37. Patel P, Hanes M, Jassal N, Pendem K, Azeem N, Deer TR. Lumbar Spine. In: *Essentials of Radiofrequency Ablation of the Spine and Joints*. 1st ed. Deer TR, Azeem N, eds. Springer Nature Switzerland AG, Cham; 2021.
38. Datta S, Lee M, Falco FJ, Bryce DA, Hayek SM. Systematic assessment of diagnostic accuracy and therapeutic utility of lumbar facet joint interventions. *Pain Physician*. 2009; 12(2):437-460.
39. Schwarzer AC, Wang SC, Bogduk N, McNaught PJ, Laurent R. Prevalence and clinical features of lumbar zygapophysial joint pain: a study in an Australian population with chronic low back pain. *Ann Rheum Dis*. 1995; 54(2):100-106. doi: 10.1136/ard.54.2.100.
40. Manchikanti L, Singh V, Pampati V, et al. Evaluation of the relative contributions of various structures in chronic low back pain. *Pain Physician*. 2001; 4(4):308-316.
41. Manchikanti L, Singh V, Pampati V, Damron KS, Beyer CD, Barnhill RC. Is there correlation of facet joint pain in lumbar and cervical spine? An evaluation of prevalence in combined chronic low back and neck pain. *Pain Physician*. 2002; 5(4):365-371.
42. Afifi A, Ringe M, Sobottke R, Oikonomidis S, Johan Teuben MP. Lumbar facet joint radiofrequency denervation therapy for chronic low back pain: enhanced outcome compared with chemical neurolysis (ethyl alcohol 95% or glycerol 20%). *Int J Spine Surg*. 2022; 16(1):33-41. doi: 10.14444/8175.
43. Li H, An J, Zhang J, et al. Comparative efficacy of radiofrequency denervation in chronic low back pain: a systematic review and network meta-analysis. *Front Surg*. 2022; 9:899538. doi: 10.3389/fsurg.2022.899538.
44. Turtle J, Miller S, Yang A, Hurley RW, Spina N, McCormick ZL. Considerations for lumbar medial branch nerve radiofrequency at spinal motion segments adjacent to a fusion construct. *Pain Med*. 2023; 24(2):165-170. doi: 10.1093/pm/pnac118.
45. Manchikanti L, Kaye AD, Soin A, et al. Comprehensive evidence-based guidelines for facet joint interventions in the management of chronic spinal pain: American Society of Interventional Pain Physicians (ASIPP) guidelines. *Pain Physician*. 2020; 23(3S):S1-S127.
46. Lo Bianco G, Misseri G, Stogicza AR, et al. Radiofrequency ablation for chronic lumbar zygapophysial joint pain using a V-shaped active tip needle: an observational retrospective study. *J Pain Res*. 2023; 16:1243-1255. doi: 10.2147/JPR.S406714.
47. Maccagnano G, Noia G, Cassano GD, et al. Thermal

- radiofrequency versus cooled radiofrequency in patients with sacroiliac joint pain: a systematic review of the literature and pooled analysis of clinical outcomes. *J Neurosurg Sci.* 2022; 66(6):485-493. doi: 10.23736/S0390-5616.22.05525-4.
48. Maalouly J, Rao PJ. Cooled radiofrequency ablation of the sacroiliac joint a retrospective case series. *BMC*

- Musculoskelet Disord.* 2023; 24(1):261. doi: 10.1186/s12891-023-06344-7.
49. Andersen GØ, Milosevic S, Jensen, et al. Coccydynia-The efficacy of available treatment options: a systematic review. *Global Spine J.* 2022; 12(7):1611-1623. doi: 10.1177/21925682211065389.