

RESEARCH ARTICLE

Developing an Appropriateness Criteria for Knee MRI Using the Rand Appropriateness Method (RAM)-2013

Hossein Ebrahimipour, MD; Seyedeh Zahra Mirfeizi, MD; Ali Vafae Najar, MD; Amir Reza Kachooei, MD; Amir Shahriar Ariamanesh, MD; Reza Ganji, MD; Habibollah Esmaeeli, MD; Hedayat Salari, MD; Marjan Vejdani, MSc

Research performed at Mashhad University of Medical Sciences, Mashhad, Iran

Received: 5 December 2013

Accepted: 4 March 2014

Abstract

Background: Knee pain is one of the most common reasons patients visit their physician. In this regard Magnetic Resonance Imaging (MRI) is the tool of preference for diagnosis. The aim of this study was to determine appropriate guidelines for knee MRI administration using the RAND Appropriateness Method (RAM)-2013

Methods: This qualitative study was done in the Mashhad University of Medical Sciences in 2013. The most appropriate approved knee MRI administration clinical guidelines were evaluated using Guidelines Evaluation and Research Appraisal (AGREE). Panel members consisting of six orthopedic and three rheumatologic doctors gave scores ranging from 1 to 9 for each scenario. The indications were grouped as appropriate, equivocal and inappropriate. Data were analyzed by descriptive statistics and SPSS ver. 18 software.

Results: Sixty-three scenarios were extracted from the guidelines and then the scenarios were evaluated in 26 indications. Thirty-two (50.79%) cases were considered appropriate, 12 (19.04%) cases uncertain and 19 (30.1%) cases inappropriate.

Conclusions: The RAND appropriateness method is helpful in identifying the opinion of stakeholders in health care systems. Moreover, making practical use of clinical guidelines can improve patients' quality of care and prevent unnecessary costs.

Key words: Guideline, Knee joint, MRI, RAND Appropriateness Method

Introduction

The knee joint is the most important weight bearing joint in the human body that is exposed to high mechanical pressure during all forms of activities such as standing, walking and even sitting, and this is due to weight, load bearing and being in a fixed position. Hence, the knee joint is commonly affected by intra- and periarticular disorders. In developing countries, such as Iran, knee injuries are very common due to the high use of motor vehicles and the increase of various sports participation without using suitable and sufficient preventive measures. These injuries can cause many problems; therefore, the correct diagnosis before any aggressive treatment is helpful in improving the final result (1).

Radiological diagnoses have helped physicians to

make early diagnosis and prevent unnecessary invasive procedures. However, medical imaging costs have increased globally and this may be due to its overuse. Among all medical imaging technologies, MRI is the most prominent and its use is predicted to increase (2-6).

In a study conducted by Lehnert and Bree, 74% of administrated CT and MRIs were reported as appropriate and 26% as inappropriate. Inappropriate imaging included MRI of the knee, back and shoulder (7). In evaluating the clinical appropriateness of an administrated MRI of the lumbar spine with the RAND appropriateness method, inappropriate, uncertain and appropriate administration was 56%, 24% and 20% respectively. The economic costs of inappropriate administration were also calculated to be 88,009,000 rial (around \$30,000 US) (8). Therefore, concern over

Corresponding Author: Marjan Vejdani, Department of Health and Management, School of Health, Daneshgah 18, School of Health, Mashhad, Khorasan Razavi, Iran.
Email: marjan_vejdani@yahoo.com



THE ONLINE VERSION OF THIS ARTICLE
ABJS.MUMS.AC.IR




Figure 1. The RAND/UCLA Appropriateness Method.

MRI prescription appropriateness has increased. Unnecessary MRI administration has a substantial financial burden for insurance companies and patients; thus, the need for an analytical tool to improve physicians' prescription ability is needed (7).

The RAND Appropriateness Method (RAM) helps developing appropriate guidelines based on available evidence and expert opinions. This method was designed by the RAND Institute and the University of California at Los Angeles and it has been used in many North American and European studies. RAM includes criteria or clinical scenarios (9). The steps of the RAM process is shown in Figure 1.

Evidence based performance has been proposed globally for some time, among which the Clinical Guidelines (CGs) is considered the best tool to apply evidences in clinical decisions. CGs transfers the available evidence into practical suggestions to help service providers (physicians) in clinical decision making (10). Clinical guidelines are systematically collected from the latest and most reliable scientific evidences that show classified ways of dealing with a patient with regard to priorities, effectiveness and cost-effectiveness.

In developing countries, there is an ongoing discussion about the fact that valid local evidences are limited and if the evidences from developed countries are going to be used, is there any need for duplication? In response to this question, the topic of guideline adoption has been introduced. Guideline adoption is a systematic view to available guidelines to find the most relevant one for our patients and then to combine it with cultural and regional requirements and available facilities of the local healthcare systems (11). RAM is used in many studies, especially to develop appropriate guidelines in surgical care as a research method (9, 12, 13). Since knee pain is one of the most common complains causing patients to visit the doctor and considering the fact that there has not been any study in Iran to evaluate knee MRI indications, this study was conducted with the aim of determining appropriate guidelines for knee MRI administration using RAM (14).

Materials and Methods

This qualitative study was performed to determine the

appropriateness of knee MRI administration on RAM.

The RAM steps are as follows:

1. Select reliable and suitable guides:

The available clinical guidelines related to knee MRI were found in the most related data bases and web sites. The search key words included: indication for knee MRI, MRI indication for knee, knee MRI guidelines, and guideline for prescribing knee MRI. The list of evaluated data bases from 2000 to 2013 is as follow: NICE, SBU, SIGN, PHAC_ASPC, HTA, AIHW, WHO, EMRO.WHO, G_N, NZGG, AHRQ, GUIDELINE, NCCN, FDA, FMH, IQWIG, ASCO, NIH, CDC, THECOCHRANELIBRARY, HAS-SANTE, and SANT.

We found nine guidelines that were assessed by two experts (a rheumatologist and an orthopedic physician) and from these, five of them were selected (9). Our panel of experts consisted of nine, and they evaluated these five guidelines using a translated and validated version of the Appraisal of Guidelines for Research & Evaluation (AGREE) while taking into consideration Iran's healthcare system (15). Each item in AGREE give scores as follows: strongly disagree, disagree, agree, and strongly agree, from 1 to 4 respectfully. Two guidelines were selected as the most appropriate scientific resource, for knee MRI indication based on the sum of these scores.

2. Selecting the panel of experts:

According to the RAND method, panel members should consist of at least nine experts, so six orthopedic physicians and three rheumatologists were selected because these experts are more involved in prescribing knee MRIs and their area of expertise is more relevant to guideline assessment (9).

3. Extraction indicators and scenarios:

Two selected guidelines for knee MRI indication were: the 2010 American College of Radiology (ACR) and the 2008 NHS Clinical Effectiveness Forum (16, 17). Finally clinical scenarios were defined based on accredited clinical guidelines and expert opinions.

4. Expert panel

The extracted scenarios from the two selected guidelines along with the scoring tool, designed by a researcher, were submitted to the experts for scoring. The researcher visited panel members separately in their work office, explained the scoring method and asked them to score scenarios based on their experience and knowledge. Each scenario had received a score from 1 to 9 (1 means that MRI administration is quite inappropriate, 5 means uncertain and 9 means appropriate).

5. Step of analyzing panel scores

Based on appropriateness scores the scenarios were divided into three groups:

Appropriate: scenarios with a mean score of 7-9 with the panel members having a majority consensus.

Uncertain: scenarios with a mean score of 4-6 that did not reach consensus agreement.

Inappropriate: scenarios with a mean score of 1-3 with total panel agreement (9).

The study was approved by Research Committee of the

Table 1. Appropriateness of MRI administration scenarios

Appropriateness	Frequency	Percent
Appropriate	32	50.7
Uncertain	12	19.04
Inappropriate	19	30.1

Mashhad University of Medical Sciences and the collected data were analyzed using descriptive statistics and SPSS ver.18 (SPSS Inc., Chicago, IL).

Results

According to RAM, in order to score the evidence, at first we conducted a meticulous literature review to find the latest available scientific evidence. The most important and related medical electronic data banks and accredited and governmental web sites were searched and nine clinical guidelines related to knee MRI were found. In the next step, based on the opinion of the expert panel head and one of the members, five clinical guidelines were chosen. Then all the panel members evaluated these five clinical guidelines based on the validated translated version of the AGREE clinical guideline evaluation tool. Based on AGREE two guidelines belonging to the American College of Radiology (ACR) and NHS Clinical Effectiveness Forum were selected (16, 17).

MRI indications of two clinical guidelines along with panel opinions for MRI administration were combined. Out of the selected guidelines 63 scenarios for knee MRI were extracted. Scenarios were evaluated in 26 indications. The scenarios' appropriateness is presented in Table 1 and several samples of the scenarios are shown in Table 2.

Discussion

The aim of this study was to identify suitable guidelines

for knee MRI administration using the RAND Appropriateness Method (RAM).

The present study is the first study to be done in Iran that used the RAND appropriateness method to design customized guidelines to prescribe knee MRI. Most of the studies that used RAM focused on medical or surgical methods by considering all or part of the following items: frequently used method, one that has high mortality and morbidity, one that used considerable resources, one that has a practice rate widely dissimilar in different geographical areas, and a controversial method regarding its use (9). Many European and US studies had used RAM in clinical cares and reported valid results. Heikes (1994), Stewart *et al* (2010), and Salari *et al's* (2013) studies evaluated RAM and concluded that it is a reliable and valid method for assessing the appropriateness of care (18).

Knee pain is one of the most common complains that cause patients referral to physicians, and on average is seen in 20% of the population (14, 19). The healthcare system's limited resources and high cost of MRI demand that physicians pay more attention in administrating MRIs. Therefore designing and performing a clinical guideline in this regard is necessary.

In addition, physicians need daily reliable information about disease prevention, diagnosis and treatment. The large amount of medical information, their differences in validity, inability of the physician to properly evaluate the information and being able to separate accredited from non-accredited information, differences in physicians' decisions about one particular disease, out of date information, inaccessibility to up to date information and the irregularities of scientific texts make the use of current information necessary (20, 21). Therefore, the goal of clinical guidelines is to inform health providers about the latest medical findings and then to change their behavior. However, this process is complex and so has to be performed systematically (10).

Table 2. Samples of MRI indications and scenarios of MRI administrations based on expert panel

Indications / Scenarios		Appropriateness
Indications (1)	Meniscal disorders	-----
Scenario (1-1)	No displaced	Appropriate
Scenario (1-2)	Displaced tears	Appropriate
Scenario (1-3)	Discoïd menisci	Appropriate
Scenario (1-4)	Meniscal cysts	Appropriate
Scenario (1-5)	Complications of meniscal surgery	Appropriate
Indications (2)	Muscle and tendon disorders	-----
Scenario (2-1)	Partial and complete tears	Appropriate
Scenario (2-2)	Tendonopathy	Appropriate
Scenario (2-3)	Ischemia	Inappropriate
Scenario (2-4)	Patellar tendon sleeve avulsions	Appropriate
Scenario (2-5)	Tears Retinacula	Appropriate
Scenario (2-6)	Non-specific inflammation of the tendon	Uncertain

Evidence based medicine is the correct, proper and wise use of the best common evidences in clinical decision making for individual patient. Its best advantage is the use of the most common evidence in clinical practice, healthcare service and management decision making (21). Early researches about the rate of using accredited scientific evidences in clinical and therapeutic decision making and the compliance of provided clinical care with accredited scientific evidence showed disappointing results. It was determined that a deep gap exists between the results of scientific researches and taken practical clinical decisions, to the extent that only 10-20% of physician interventions were in compliance with scientific evidence (22).

Nejat *et al's* study, evaluating the information resources used by general practitioners in Tehran for acquiring knowledge, showed that less than 32% of participants were familiar with clinical guidelines. Despite the positive attitude of physicians toward clinical guidelines and considering them as useful and trustable, they believed that many barriers exist for using them. Moreover, the low familiarity of physicians with clinical guidelines and their low use show the problems in providing customized evidences and developing customized clinical guidelines (23).

Since defined scenarios in this study are a combination of international scientific literature and the experience of national experts, the developed guideline is accredited and adapted and can help physicians in proper decision making.

Study limitation

Since this was a qualitative research and considering the fact that by changing panel members we could have different results, the study results should be generalized with caution.

Conclusion

The RAND appropriateness method is very useful in identifying the opinion of stakeholders in systems with limited resources. Moreover, developing, customizing and performing clinical guidelines are a challenge in the Iranian healthcare system. Hence, making clinical guidelines practical can lead to the improvement of quality of care in the healthcare system. The result of this study can be used for developing national guidelines and aid health policy makers to consider adopted national clinical evidence in decision making.

Acknowledgements

This article is the result of a Master thesis in Health Care Management, School of Health, Mashhad University of Medical Sciences, (grant number: 911105), approved by the university research chancellor. The researcher appreciates the cooperation of Dr. Arash Rashidian, Dr. Ali Moradi, Dr. Maryam Sahebari, and Dr. Vahid Azhdari.

Hossein Ebrahimipour MD

Health Sciences Research Center, Department of Health and Management, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

Seyedeh Zahra Mirfeizi MD

Rheumatic Diseases Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

Ali Vafaei Najar MD

Health Sciences Research Center, Department of Health and Management, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

Amir Reza Kachooei MD

Orthopedic Research Center, Mashhad University Of Medical Sciences, Mashhad, Iran

Amir Shahriar Ariamanesh MD

Mashhad University of Medical Sciences, Mashhad, Iran

Reza Ganji MD

Department of Orthopedic Surgery, North Khorasan University of Medical Sciences, Bojnurd, Iran

Habibollah Esmaeeli MD

Health Sciences Research Center, Department of Biostatistics, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

Hedayat Salari MD

Department of Health Management and Economics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

Marjan Vejdani MSc

Student Research Committee, School of Health, Mashhad University of Medical Sciences, Mashhad, Iran

References

- Habashizadeh T, Heravi MT, Makhmalbaf H, Esmaeeli H. Evaluation of MRI results in clinical suspected ACL tears and its comparison with arthroscopic findings in 100 patients approaching the Ghaem Hospital from 2005 up to 2006. Mashhad: Mashhad University of Medical Sciences; 2007.
- Greenberg D, Peterburg Y, Vekstein D, Pliskin JS. Decisions to adopt new technologies at the hospital level: insights from Israeli medical centers. *Int J Technol Assess Health Care*. 2005;21(2):219-27.
- Fuchs VR, Sox HC Jr. Physicians' views of the relative importance of thirty medical innovations. *Health Aff (Millwood)*. 2001;20(5):30-42.
- Passariello R. Cost containment and diffusion of MRI: oil and water?. The situation in Europe. *Eur Radiol*. 1997;7:259-62.

5. Hillman AL, Schwartz JS. The diffusion of MRI: patterns of siting and ownership in an era of changing incentives. *AJR Am J Roentgenol.* 1986;146(5):963-9.
6. Oh EH, Imanaka Y, Evans E. Determinants of the diffusion of computed tomography and magnetic resonance imaging. *Int J Technol Assess Health Care.* 2005;21(1):73-80.
7. Lehnert BE, Bree RL. Analysis of appropriateness of outpatient CT and MRI referred from primary care clinics at an academic medical center: how critical is the need for improved decision support? *J Am Coll Radiol.* 2010;7(3):192-7.
8. Salari H, Ostovar R, Esfandiari A, Keshtkaran A, Sari AA, Manesh HY, et al. Evidence for Policy Making: Clinical Appropriateness Study of Lumbar Spine MRI Prescriptions Using RAND Appropriateness Method. *Int J Health Policy Manag.* 2013;1(1):17-21.
9. Fitch K, Bernstein SJ, Aguilar MD, Burnand B, LaCalle JR, Lázaro P, et al. The RAND/UCLA Appropriateness Method User's Manual; 2001.
10. Ashorkhani M, Gholami Z, Majdzade SR, Maleki K, Nejat S, Nejat S, et al. EBM. Department of Knowledge Translation; 2009.
11. Effectiveness. Bushehr university of medical sciences 2012; Available from: <http://www.bpums.ac.ir/fa/DynPages-1156.htm>.
12. Nicollier-Fahrni A, Vader JP, Froehlich F, Gonvers JJ, Burnand B. Development of appropriateness criteria for colonoscopy: comparison between a standardized expert panel and an evidence-based medicine approach. *Int J Qual Health Care.* 2003;15(1):15-22.
13. Porchet F, Vader JP, Larequi-Lauber T, Costanza MC, Burnand B, Dubois RW. The assessment of appropriate indications for laminectomy. *J Bone Joint Surg Br.* 1999;81(2):234-9.
14. Rezaee Moghadam F, Azma K, Raeesolsadat SA, Saadati N, Shamsodini A, Naseh I. Effect of basic military training on knee pain and flexibility of the muscles surrounding. *Journal of Military Medicine.* 2006;8(4):260-55.
15. Irani S, Rashidian A, Yousefi-Nooraie R, Soltani A. Evaluating clinical practice guidelines developed for the management of thyroid nodules and thyroid cancers and assessing the reliability and validity of the AGREE instrument. *J Eval Clin Pract.* 2011;17(4):729-36.
16. Priorities and clinical effectiveness forum. MRI and internal knee derangement guideline. 2008; Available at: http://www.derbyshire.nhs.uk/referrals/mri_and_internal_knee_derangement_guideline_2005.pdf
17. The American College of Radiology. ACR-SSR practice guideline for the performance and interpretation of magnetic resonance imaging (MRI) of the knee. 2010; Available at: <http://www.acr.org/Quality-Safety/Standards-Guidelines/Practice-Guidelines-by-Modality/MRI>
18. Salari H, Esfandiari A, Ostovar R, Keshtkaran A. Developing Appropriateness criteria for MRI of Low BackPain(LBP)using Rand Appropriateness Method (RAM). *Armaghane-danesh, Yasuj University of Medical Sciences Journal (YUMSJ).* 2013;18(2).
19. Jackson JL, O'Malley PG, Kroenke K. Evaluation of acute knee pain in primary care. *Ann Intern Med.* 2003;139(7):575-88.
20. Soltani A, Allameh F, Hoseini M, Froghi N, Sichani BS. Evidence-Based Medicine. *Journal of Science and Technology.* 2007;1(3).
21. Jedi FR, Moraveji A, Abazari F. Awareness and use of evidence-based medicine among general practitioners, specialists and residents of areas of the country. *Urmia Medical Journal.* 2012;23(6).
22. Kachoei MA, Sdeghizadeh A, Taheri H, Nezhad NK, Pour SM, Kabiri P, et al. Does our clinical decision-making is evidence-based? *Jornal of Isfahan Medical School.* 2010;28(121).
23. Nejat S, Gholami Z, Maleki K, Ashorkhani M, Ahghari S, Nejat S, et al. Sources of information used for the knowledge gained by physicians in Tehran. *Secretariat of Research and Policy Development;* 2009.