

REVIEW ARTICLE

Surgical Management of Hip Problems in Myelomeningocele: A Review Article

Taghi Baghdadi, MD; Reza abdi, MD; Ramin Zargar Bashi, MD; Hossein Aslani, MD

*Research performed at Emam Reza hospital, Birjand University of medical science, Birjand, Iran**Received: 10 July 2014**Accepted: 20 February 2016***Abstract**

Background: Children with myelomeningocele (MMC) develop a wide variety of hip deformities such as muscle imbalance, contracture, subluxation, and dislocation. Various methods and indications have been introduced for treatment of muscle imbalances and other hip problems in patients with MMC but there is no study or meta-analysis to compare the results and complications. This review aims to find the most acceptable approach to hip problems in patients with MMC.

Methods: MEDLINE was searched up to April 2015. All study designs that reported on the outcomes of hip problems in MMC were included. From 270 screened citations, 55 were strictly focused on hip problem in MMC were selected and reviewed.

Results: Complex osseous and soft tissue reconstructive procedures to correct hip dysplasia and muscle balancing around the hip are rarely indicated for MMC patients without good quadriceps power.

Conclusion: Over the years a consensus on the best algorithm for treatment of hip dislocation in myelomeningocele has been missing, however, muscular balancing with/without osseous procedure seems a reasonable approach especially in unilateral mid-lumbar MMC.

Keywords: Hip dislocation, Hip dysplasia, Myelomeningocele, Meningomyelocele, Spina bifida, Teratology

Introduction

Myelomeningocele (MMC) is a saclike structure containing cerebrospinal fluid and neural tissue, caused by a failure of the neural tube to close during the fourth week of gestation (1). It is a common malformation, occurring in approximately 1 in 1,000 live births. The cause is unknown, but, both genetic and environmental factors have significant roles. MMC may be located anywhere along the neuraxis, but, 75% occur in the lumbosacral region (2). In the 1960s, effective techniques were developed for shunting hydrocephalus and early closure of neural tube defects. As a result, orthopaedic surgeons were presented with the challenge of managing a population of children who had MMC (3). Initially, the musculoskeletal problems in these children were treated with the modalities that had been learned from the treatment of poliomyelitis. However, it soon became apparent that the management of children who have MMC was not so simple because of additional factors such as decreased sensation of the lower extremities or encephalopathy that impair

coordination and results in the loss of strength of the lower and upper extremities and muscle imbalance that affects skeletal development over the entire period of growth (1-3). Progressive neurological deterioration may occur in MMC because of tethered cord syndrome, syringomyelia or hydrocephalus (4, 5). Hydrocephalus in association with a type II Arnold-Chiari defect is common and develops in 80% of children with MMC but only 30% of patients need a neurosurgical shunt (2, 6). Ambulation in patients with MMC is affected not only by the neurological level of motor performance, but also by other factors such as age, obesity, cognitive status, motivation, spasticity, upper limb functional status, orthopedic deformities and intelligence (7, 8). At minimum follow-up of 20 years 42% of MMC had normal IQ but only 8% achieved college degrees (6).

Materials and Methods

We reviewed articles concerning different treatments of hip problems in MMC. The search engine was MedLine (PubMed) and the keywords used were:

Corresponding Author: Reza Abdi, Emam Reza Hospital, Birjand University of Medical Science, Birjand, Iran
Email: reza1352abdi@gmail.com



THE ONLINE VERSION OF THIS ARTICLE
ABJS.MUMS.AC.IR

meningomyelocele + hip; myelomeningocele + hip; and spina bifida + hip.

Until April 2015, 270 articles were retrieved for review. Articles were included if they provided an abstract and were written in English. Of those, 55 were selected and reviewed when they were strictly focused on the topic of this article. In addition, references of these articles were examined to ensure no compelling literature had been overlooked. The types of studies reported have a low level of evidence (most of them are level III-IV studies).

Results

Classification

The neurological level of lesion is an essential factor influencing ambulation in children with MMC and hip problem (9). The most frequently used classification systems of the neurological level of lesions in the literature are Sharrard (1964), Hoffer (1973), Lindseth (1976), Ferrari (1985), McDonald (1991), and Broughton (1993)(7).

Hoffer and colleagues' classification was based on studies of 56 patients with MMC aged between 5 and 42 years. The resulting classification system has four categories: thoracic, upper lumbar, lower lumbar, and sacral. Also they considered four levels of functional ambulation consisting of community ambulators, household ambulators, non-functional ambulators and non-ambulators (7, 8). In 1999, Bartoneck analyzed these six commonly used classification system in patients with MMC and suggested that it is not possible to compare neurological levels in different classification system to each other or definitely correlate the results to the functional ambulation levels of Hoffer classification (7, 8).

A modification of the classification system described by Asher and Olson based on the lowest level of antigravity (at least grade-3) muscle strength on the patient's best side to define the neurological level is simple to use, and has been helpful in predicting gross motor function and potential problems (3, 10).

Thoracic level: Patients who have only functional thoracic roots may be able to walk during the first decade of life, but, they become dependent on a wheelchair as they attain adult body mass. They do not have active movement of the lower extremity muscles and have some abduction, external rotation, and flexion contracture around their hip (11-14).

Upper lumbar level (L1, L2): They have active flexion of the hip but seldom retain the capacity for functional walking after reaching the adulthood. A contracture of the unopposed hip flexors typically develops in these patients. A mild contracture of the unopposed hip adductors may also occur, however, restriction in hip abduction is usually mild and is not clinically important(3, 12).

Mid-lumbar level (L3, L4): Patients who have mid-lumbar MMC typically have normal strength in the hip flexors and adductors but no function in hip extensors or abductors. Therefore, a flexion contracture of the hip and some limitation of abduction frequently develop in these patients. More importantly, this pattern of muscle

imbalance predisposes to progressive subluxation of the hip (8).

Lumbosacral level: The probability of subluxation or a severe flexion contracture of the hip is low in patients who have lumbosacral MMC. This is particularly true if the MMC is at the sacral level, but adequate stability and an adequate range of motion of the hip are usually maintained even in patients with fifth lumbar MMC who have only grade 2 or 3 strength of the hip abductors and absent or trace strength of the hip extensors (15, 16). Apparently, this degree of activity of the hip abductors combined with activity of the hamstrings can be an effective counterbalance to hip flexors and adductors of normal strength. These patients should have periodic radiographs during early childhood to monitor the development of the hip joint. It was previously thought that all patients who had sacral MMC were able to walk independently; However, in a study, among 36 patients who were evaluated as adults, 6 had become dependent on a wheelchair as a result of neurological deterioration, ulceration of the feet, and other problems (15).

Natural history: Neurological level is a critical indicator in determining the ambulation capacity, functional capability and hip deformity. 56% of adults with MMC are unemployed and 43% use wheelchair (6). All patients in L2 level are wheelchair dependent and 60% of patients bellow L2 level use wheelchair sometimes. Maximal level of ambulation will achieve at age four to six years old and if the child cannot stand at six years old then walking will be impossible (16).

Samuelsson *et al.* studied factors determining ambulation in 163 patients with MMC by a multivariate statistical method and concluded that severe scoliosis was closely, age was moderately, and hip flexion contracture was slightly related to the inability to walk, while pelvic obliquity, hip dislocation, or knee flexion contracture were not (17). Also several other studies have demonstrated that the ability to walk is not affected by dislocation of the hip in patients who have thoracic or upper lumbar MMC (11-14, 18-21). Therefore, complex osseous and soft tissue reconstructive procedures to correct hip dysplasia and muscle balancing around the hip are rarely indicated for these patients since relocation of the hip with a good radiograph does not mean functional advantage and may cause complications, such as pathological fracture or devastating stiffness of the hip joint (6, 22). Although

Table 1. Asher and Olson classification of MMC

Level	Function
Thoracic	No grade-3 strength in muscles of lower limbs
L1-L2	Hip flexion or Adduction
L3	Knee extension
L4	Knee flexion
L5	Ankle dorsiflexion
S1	Ankle plantar flexion

Patients with MMC can also be functionally classified in four groups: (3)

based on opinion of some well-respected authors, surgical reduction of paralytic hip dislocations in ambulatory MMC patients is costly and offers little obvious benefit but many authors believe that it is appropriate in low-level unilateral dislocations and it seems that, in the large series, Sharrard procedure was done virtually on all patients who had mid-lumbar MMC (19-25-26, 28-35). In 1969 Menelaous believed that dislocated hip in MMC should always be reduced unless there is severe paralysis or low intelligence, but, in 1984 he said: "We feel that surgery of this magnitude of iliopsoas (ILP) transfer should be performed only on those patients likely to walk in AFO into adult life" (3, 11, 12).

Up to 80% patient with MMC have a problem or weakness in upper extremity and Charney found that, with bracing and gait training, 52% of 87 patients who had MMC at upper level were able to walk about the community at age five but deficient balance reaction and weakness of the upper extremities coupled with the extensive bracing that is needed, prevents some of these children from achieving a functional walking ability (16, 36). Children who have upper level MMC rarely retain the ability to walk after reaching the adulthood and hip dislocation in MMC is not painful compared to CP, so, some authors have advocated an intensive program of bracing and gait training during early childhood. The potential benefits of walking by patients with MMC at upper level include strengthening of the upper extremities, protection against obesity, and prevention of contractures of the lower extremities. However, the most important benefit is the psychological boost and tremendous sense of accomplishment that these children express when they achieve the ability to be upright and to move around a room like other children of their age (37, 38). Furthermore, in a study, patients with this level of MMC who were managed with bracing and early walking had fewer fractures and pressure sores and were more independent in transfers, even when they eventually switched to a wheelchair, compared with patients who had always used a wheelchair. However, these children were hospitalized more often, for operative procedures to allow bracing (3, 38).

Muscle balancing procedures: In 1952 Mustard introduced lateral transfer of the ILP through the anterior iliac window to the greater trochanter, but Mustard's procedure was abandoned because it did not reinforced hip extensors. The story of the surgical treatment of muscle imbalance around the hip in MMC began by Sharrard (1964). He introduced the posterolateral transfer of iliopsoas to the greater trochanter (39, 40). Also, he found no hip dislocations or flexion deformities in limbs without any innervation below T12 and no active muscles around the hip. He said that if there is no muscle activity around the hip then hip dislocation does not occur; however, nowadays we know that this is not correct (41). In a multicenter study of 1061 patients with MMC, measurement of the flexion contractures of the hip in older children (9-11 years old) revealed that the greatest average value of flexion contracture was in the patients who had thoracic or upper lumbar MMC. Dislocation tended to occur by the age of three to four

years in the patients who had mid-lumbar MMC, but those who had thoracic or upper lumbar MMC continued to have dislocation of the hip even after the age of ten years (18).

Time of transfer: Posterolateral transfer of the ILP should be limited to a selected group of patients. Sharrard reported some deformities of patients, presented at the age of one year after birth in 183 children with MMC, so, in 1983 he recommended the transfer should be done before development of osseous deformity (between one and two years of age), combined with an adductor release, and should be limited to patients who have fourth lumbar MMC (29). Also later two studies demonstrated the Sharrard procedure or external oblique muscle transfer before age one is unsuccessful. In the study by Stillwell and Menelaus, five out of nine patients with ILP transfer before one year old, were not able to walk during the follow-up, at least ten years postoperatively but they provided useful data concerning the effectiveness of the Sharrard procedure in other ages (30). Also, Tosi *et al.* transferred external oblique muscle and femoral osteotomy after initial treatment with a Pavlik Harness and reported redislocation of two of four hips. The reason for the redislocation was unclear, but the early operation, or early wrong grading of quadriceps may explain them (31). Because early grading of quadriceps power in first 3 years of life give a reliable assessment of future ambulatory ability, but it should be noted that early grading of quadriceps power compared to grading at later ages, only 56 out of 109 assessments remained the same (42).

Absence of flexor power after ILP transfer and role of flexion contracture around the hip: Stillwell and Menelaus have reported 47 patients with ILP transfer and adductor release to obtain 60 degrees of abduction more than 10 years ago among which, 32 (68%) were community walkers, 3 were household walkers and 12 were non-walkers. Compared to other published reports, these patients did not lose their walking ability that could be jeopardized by the loss of hip ILP flexor power. Furthermore, all except three of the community walkers were able to climb and descend stairs after ILP transfer (30).

Many patients with intact flexor and absent extensor function around the hip developed hip flexion contractures. Contracture in household ambulatory patients ranged from 0 to 45 (average 23) degrees. As hip flexion contracture did not appear to impair the ambulatory ability when the reciprocating gait orthosis was used, the surgical release of functioning hip flexor musculature if the use of this device is being considered, is not indicated and up to 30 degree of the hip flexion contracture is acceptable (16, 41, 43).

Shurtleff *et al.* analyzed 5,147 serial measurements of the range of hip extension in 966 patients with spina bifida and concluded that the contractures were generally present in the first few months of life (physiologic flexion posture); this then diminished during the first 27 months in all but those with thoracic lesions. They demonstrated that the contractures reappear or worsen between the ages of 3 and 6 years and are not merely due

to muscle imbalance, sitting posture, or these factors in combination. Hence, surgical management is seldom appropriate until after that age (44). Also Frowley showed that successful isolated anterior hip release is not related to the neurological level or the operation age; also, recurrence of contracture correlated with the walking ability of the child at the latest follow-up (43).

Muscle balancing procedure should be done in located hip or subluxed hip? Based on the fact that muscular imbalance of the hip is responsible for progressive hip dislocation and secondary osseous deformity, Weisl *et al.* (1988) reported ILP transfer on 54 hips, 23 of them (42.6%) had located hips at the moment of the operation (32). Molloy (1986) combined the ILP transfer with femoral varus derotation osteotomy on 26 hips; five hips (19.2%) were located and did not have dysplastic changes (33). Bunch and Hakala (1984) performed an ILP transfer on 32 hips, but three of them (9.4%) showed no abnormal changes (28). Lorente (2005) performed the ILP transfer on 24 hips, 41.4% of hips were located in his series (45). However, it is now understood that progressive dysplasia does not develop in all patients who have MMC at the mid-lumbar level, and experienced observers, such as Broughton and Menelaus, after a review of 1061 patients, concluded muscular imbalance is not a significant factor in the production of flexion deformity or dislocation of the hip and they do not recommend prophylactic surgery. They recommend a selective approach for operative intervention only for patients in whom subluxation has developed when have at least grade-4 strength of the quadriceps (3, 41). Although Lorente (2005) concluded that ILP transfer procedure has value in obtaining hip stability and walking ability in carefully selected MMC with L3 paralysis but it is unclear whether prophylactic posterolateral transfer ILP should be done in located hip at L3 level hip or should be done only in hips that are going to be sublux (45).

ILP or external oblique transfer: It is now clear that posterolateral transfer of the ILP does not provide active extension or abduction against gravity, and it is doubtful that this out-of-phase transfer provides any noticeable extension or abduction during the gait cycle (3). The transferred ILP or external oblique muscle is unlikely to provide significant active abduction or extension power, however ILP transfer may have a beneficial function as a tenodesis (31, 46). In our limited experience some abduction power is achieved after ILP transfer only in supine position. A recent study using three dimensional gait analysis revealed no improvement in abnormal pelvic obliquity in patients with fourth lumbar MMC who had been managed with ILP transfer (47). Transfer of the external oblique muscle has been advocated as an alternative to transfer of the iliopsoas for patients who have MMC at the mid-lumbar level as well as a dysplastic hip (31, 34, 48). This procedure does not weaken the iliopsoas; therefore, the power of the hip flexors and the ability to climb stairs should be maintained. Some authors have stated that transfer of the external oblique muscle improves hip mechanics during mid-stance, however, gait-analysis studies have demonstrated that

the transferred external oblique muscle mainly functions during the swing phase of gait therefore it is doubtful that this transferred muscle simulates the activity of the hip abductors or extensors during walking (35). In a comparative study external oblique muscle transfer did not provide a clinically important improvement in functional recovery in patients with L3 to L5 level when added to periarticular release of contractures and bony procedures (49). In spite of these articles it is difficult to compare the results of ILP and external oblique muscle transfer because a comparative study is not yet done between two transfers in MMC and there is not any general agreement on the choice of the muscle transfer (30, 31, 34).

Triple transfer: Phillips and Lindseth (1992) described the results of triple transfer of the external oblique muscle to the greater trochanter, the hip adductors to the ischium and the tensor fasciae latae to a more lateral position on 89 hips (34). Although those authors reported functional walking by all patients, the duration of follow-up was not enough.

Transfer in bilateral dislocation: According to Menelaus (1980) and Carroll (1987), due to leg length discrepancy, pelvic obliquity develop, resulting pressure sores, poor sitting balance, and a negative influence on the status of spinal column, bilateral dislocations should be treated only when full power in quadriceps muscle exists and unilateral dislocation should be treated regardless of the level of paralysis (50), but Robert (1994), Crandall (1989), and Sherk (1991) have reported that the incidence of these problems with the exception of a leg length discrepancy is not significantly influenced by unilateral dislocation of hip and they do not recommend the procedure in inappropriate set (19).

The role of osseous procedure in combination with muscular balancing procedure: Muscle transfers for the treatment of dysplasia of the hip associated with MMC are insufficient to correct severe osseous deformities. Misalignment of the proximal part of the femur or the acetabulum is particularly common after the age of three years and this problem should be corrected either before or at the same time as the muscle transfers (3). A femoral varus derotation osteotomy corrects abnormal valgus angulation and anteversion. The type of pelvic osteotomy that best serves these patients is less clear. These patients often have global acetabular deficiency (51). For that reason, a Pemberton or modified Dega procedure may be better than other pelvic osteotomies that are commonly used for the management of young children who have CDH (3). Chiari osteotomy did not achieve long-term hip stability and good results in many patients (52, 53). In a study of 34 children (66 hips) with third, fourth, or fifth lumbar MMC who had a femoral osteotomy combined with transfer of the external oblique and adductor muscles, Tosi *et al.* reported the maintenance of stability of 37 (73%) of 51 hips in the 26 children who remained neurologically stable; however, only eight of 15 hips in children who had progressive loss of neurological function remained stable (31). The poorest results were for the hips that had dislocated previously. Only two out of 10 hips in this group had

successful results. The average duration of follow-up in that study was relatively long (10.9 years), but the wide range of follow-up (0.7 to 20.0 years) limits conclusions concerning about functional status when those children reached adult body sizes. At the most recent evaluation, 21 (81 per cent) of 26 children who did not lost neurological function during follow-up were able to walk about the community (31). So even after osseous procedure in MMC, progressive loss of neurological function is an important factor for the stability of hip and ambulation. The combination of one stage pelvic and femoral osteotomy and transiliac psoas transfer can be effective in selected patients with MMC (33).

Discussion

Treatment in the first year of life: As mentioned above, muscle transfer has poor outcomes before the age of one year, so, treatment of subluxation or dislocation of the hip in midlumbar MMC is difficult during the first year after birth. Typically, the problem is noted in the first few months of life. Use of a Pavlik Harness or some other brace designed for CDH is seldom successful over the long term for these patients (3). Furthermore, these braces exacerbate a flexion contracture in an infant when the hip extensors are nonfunctional (3). Based on Green (1998) and Breed (1982), when dysplasia develops in the first year of the life in patients who have midlumbar MMC, it develops by the age of three or four months and many of these patients probably have dislocation of the hip at birth, but the treatment of other medical problems prevents its documentation. If intervention is delayed, the rapid growth during infancy coupled with an underlying muscle imbalance may result in severe dysplasia that cannot be stabilized with soft-tissue procedures. Because pavlik can induces flexion contracture of hip and is not successful to treat muscle imbalance, they recommend an ILP recession and adductor myotomy (3, 54). They defined the "bowstring" force of the iliopsoas, the force applied to the femoral head as the tendon angles across the hip joint, which they believe is an important cause of dislocation of the hip in patients with a midlumbar MMC. An operative procedure consisting of ILP recession and suture of its tendon to the anterolateral hip joint capsule

has been developed and used in 10 dislocated and 9 subluxated hips. Secondary femoral varus derotation osteotomy for valgus and anteversion was performed on five hips with subluxation. Finally 16 out of 19 hips were stable, two have subluxation, and one was dislocatable. They recommended early surgical treatment to prevent secondary adaptive changes in the hip (54).

Treatment of progressive subluxation of the hip in adolescent patient: Progressive subluxation of the hip in adolescent patient with midlumbar MMC presents a dilemma. Muscle imbalance coupled with complicated hip dysplasia leads to a substantial rate of recurrent subluxation. There is no answer for this group of patients. Certainly, the evaluation should include assessment for a possible tethered cord syndrome and consideration of a CT scan with reconstruction to define the extent and location of the acetabular deformity (51, 55). Base on Instructional Course Lectures AAOS (1998), the treatment must be individualized (3). Green have most often performed a Chiari pelvic osteotomy and femoral varus derotation osteotomy in this group of patients. He have also recommended observation or no treatment for adolescent patients who have a history of operations, no functional abductor muscles, and a markedly dysplastic acetabulum. In this situation, the chance of success is low and reconstructive operations may cause the hip to become stiff and painful (3). However Mannor (1996) and Zenios (2012) showed Chiari osteotomy did not achieve long-term hip stability and good results in MMC (52, 53).

Taghi Baghdadi MD

Reza abdi MD

Emam Educational Hospital, Tehran University of Medical Science, Tehran, Iran

Ramin Zargar Bashi MD

Children's Medical Center, Tehran University of Medical Science, Tehran, Iran

Hossein Aslani MD

Shohada Educational Hospital, Golshahr Town, Tabriz, Iran

References

1. Feeley BT, Ip TC, Otsuka NY. Skeletal maturity in myelomeningocele. *J Pediatr Orthop.* 2003; 23(6):718-21.
2. Stein SC, Schut L. Hydrocephalus in myelomeningocele. *Childs Brain.* 1979; 5(4):413-9.
3. Greene WB. Treatment of hip and knee problems in myelomeningocele. *Instr Course Lect.* 1999; 48:563-74.
4. Mazur JM, Stillwell A, Menelaus M. The significance of spasticity in the upper and lower limbs in myelomeningocele. *J Bone Joint Surg Br.* 1986; 68(2):213-7.
5. Mazur JM, Menelaus MB, Hudson I, Stillwell A. Hand function in patients with spina bifida cystica. *J Pediatr Orthop.* 1986; 6(4):442-7.
6. Roach JW, Short BF, Saltzman HM. Adult consequences of spina bifida: a cohort study. *Clin Orthop Relat Res.* 2011; 469(5):1246-52.
7. Bartonek A, Saraste H, Knutson LM. Comparison of different systems to classify the neurological level of lesion in patients with myelomeningocele. *Dev Med Child Neurol.* 1999; 41(12):796-805.
8. Hoffer MM, Feiwel E, Perry R, Perry J, Bonnett C. Functional ambulation in patients with

- myelomeningocele. *J Bone Joint Surg Am.* 1973; 55(1):137-48.
9. Fraser RK, Hoffman EB, Sparks LT, Buccimazza SS. The unstable hip and mid-lumbar myelomeningocele. *J Bone Joint Surg Br.* 1992; 74(1):143-6.
 10. Asher M, Olson J. Factors affecting the ambulatory status of patients with spina bifida cystica. *J Bone Joint Surg Am.* 1983; 65(3):350-6.
 11. Barden GA, Meyer LC, Stelling FH 3rd. Myelodysplastics--fate of those followed for twenty years or more. *J Bone Joint Surg Am.* 1975; 57(5):643-7.
 12. Bazih J, Gross RH. Hip surgery in the lumbar level myelomeningocele patient. *J Pediatr Orthop.* 1981; 1(4):405-11.
 13. De Souza LJ, Carroll N. Ambulation of the braced myelomeningocele patient. *J Bone Joint Surg Am.* 1976; 58(8):1112-8.
 14. Feiwell E. Surgery of the hip in myelomeningocele as related to adult goals. *Clin Orthop Relat Res.* 1980; 148:87-93.
 15. Brinker MR, Rosenfeld SR, Feiwell E, Granger SP, Mitchell DC, Rice JC. Myelomeningocele at the sacral level. Long-term outcomes in adults. *J Bone Joint Surg Am.* 1994; 76(9):1293-300.
 16. Canale ST, Beaty JH. *Campbell's operative orthopaedics.* 12th ed. Philadelphia: Elsevier Health Sciences; 2013.
 17. Samuelsson L, Skoog M. Ambulation in patients with myelomeningocele: a multivariate statistical analysis. *J Pediatr Orthop.* 1988; 8(5):569-75.
 18. Broughton NS, Menelaus MB, Cole WG, Shurtleff DB. The natural history of hip deformity in myelomeningocele. *J Bone Joint Surg Br.* 1993; 75(5):760-3.
 19. Fraser RK, Bourke HM, Broughton NS, Menelaus MB. Unilateral dislocation of the hip in spina bifida. A long-term follow-up. *J Bone Joint Surg Br.* 1995; 77(4):615-9.
 20. Keggi JM, Banta JV, Walton C. The myelodysplastic hip and scoliosis. *Dev Med Child Neurol.* 1992; 34(3):240-6.
 21. Lee EH, Carroll NC. Hip stability and ambulatory status in myelomeningocele. *J Pediatr Orthop.* 1985; 5(5):522-7.
 22. Canale G, Scarsi M, Mastragostino S. Hip deformity and dislocation in spina bifida. *Ital J Orthop Traumatol.* 1992; 18(2):155-65.
 23. Sherk HH, Uppal GS, Lane G, Melchionni J. Treatment versus non-treatment of hip dislocations in ambulatory patients with myelomeningocele. *Dev Med Child Neurol.* 1991; 33(6):491-4.
 24. Swaroop VT, Dias LS. Strategies of hip management in myelomeningocele: to do or not to do. *Hip Int.* 2009; 19(Suppl 6):S53-5.
 25. Sherk HH, Melchionne J, Smith R. The natural history of hip dislocations in ambulatory myelomeningoceles. *Z Kinderchir.* 1987; 42(Suppl 1):48-9.
 26. Wright JG. Hip and spine surgery is of questionable value in spina bifida: an evidence-based review. *Clin Orthop Relat Res.* 2011; 469(5):1258-64.
 27. Szulk A, Tomaszewski M, Koch A, Kotwicki T. Current experience in the treatment of neurogenic deformities of the hip joint in patients with myelomeningocele. *Ortop Traumatol Rehabil.* 2011; 13(2):125-43.
 28. Bunch WH, Hakala MW. Iliopsoas transfers in children with myelomeningocele. *J Bone Joint Surg Am.* 1984; 66(2):224-7.
 29. Sharrard WJ. Management of paralytic subluxation and dislocation of the hip in myelomeningocele. *Dev Med Child Neurol.* 1983; 25(3):374-6.
 30. Stillwell A, Menelaus MB. Walking ability after transplantation of the iliopsoas. A long-term follow-up. *J Bone Joint Surg Br.* 1984; 66(5):656-9.
 31. Tosi LL, Buck BD, Nason SS, McKay DW. Dislocation of hip in myelomeningocele. The McKay hip stabilization. *J Bone Joint Surg Am.* 1996; 78(5):664-73.
 32. Weisl H, Fairclough JA, Jones DG. Stabilisation of the hip in myelomeningocele. Comparison of posterior iliopsoas transfer and varus-rotation osteotomy. *J Bone Joint Surg Br.* 1988; 70(1):29-33.
 33. Molloy MK. The unstable paralytic hip: treatment by combined pelvic and femoral osteotomy and transiliac psoas transfer. *J Pediatr Orthop.* 1986; 6(5):533-8.
 34. Phillips DP, Lindseth RE. Ambulation after transfer of adductors, external oblique, and tensor fascia lata in myelomeningocele. *J Pediatr Orthop.* 1992; 12(6):712-7.
 35. Carroll NC, Sharrard WJ. Long-term follow-up of posterior iliopsoas transplantation for paralytic dislocation of the hip. *J Bone Joint Surg Am.* 1972; 54(3):551-60.
 36. Charney EB, Melchionni JB, Smith DR. Community ambulation by children with myelomeningocele and high-level paralysis. *J Pediatr Orthop.* 1991; 11(5):579-82.
 37. Liptak GS, Shurtleff DB, Bloss JW, Baltus-Hebert E, Manitta P. Mobility aids for children with high-level myelomeningocele: parapodium versus wheelchair. *Dev Med Child Neurol.* 1992; 34(9):787-96.

38. Mazur JM, Shurtleff D, Menelaus M, Colliver J. Orthopaedic management of high-level spina bifida. Early walking compared with early use of a wheelchair. *J Bone Joint Surg Am.* 1989; 71(1):56-61.
39. Mustard WT. A follow-up study of iliopsoas transfer for hip instability. *J Bone Joint Surg Br.* 1959; 41-B(2):289-98.
40. Mustard WT. Iliopsoas transfer for weakness of the hip abductors; a preliminary report. *J Bone Joint Surg Am.* 1952; 24 A(3):647-50.
41. Yngve DA, Douglas R, Roberts JM. The reciprocating gait orthosis in myelomeningocele. *J Pediatr Orthop.* 1984; 4(3):304-10.
42. Schopler SA, Menelaus MB. Significance of the strength of the quadriceps muscles in children with myelomeningocele. *J Pediatr Orthop.* 1987; 7(5):507-12.
43. Frawley PA, Broughton NS, Menelaus MB. Anterior release for fixed flexion deformity of the hip in spina bifida. *J Bone Joint Surg Br.* 1996; 78(2):299-302.
44. Shurtleff DB, Menelaus MB, Staheli LT, Chew DE, Lamers JY, Stillwell A, et al. Natural history of flexion deformity of the hip in myelodysplasia. *J Pediatr Orthop.* 1986; 6(6):666-73.
45. Lorente Molto FJ, Martinez Garrido I. Retrospective review of L3 myelomeningocele in three age groups: should posterolateral iliopsoas transfer still be indicated to stabilize the hip? *J Pediatr Orthop B.* 2005; 14(3):177-84.
46. Rueda J, Carroll NC. Hip instability in patients with myelomeningocele. *J Bone Joint Surg Br.* 1972; 54(3):422-31.
47. Duffy CM, Hill AE, Cosgrove AP, Corry IS, Mollan RA, Graham HK. Three-dimensional gait analysis in spina bifida. *J Pediatr Orthop.* 1996; 16(6):786-91.
48. Dias LS. Hip deformities in myelomeningocele: In Instructional Course Lectures. 40. Park Ridge, Illinois: American Academy of Orthopaedic Surgeons; 1991. p. 280-6.
49. Yildirim T, Gursu S, Bayhan IA, Sofu H, Bursali A. Surgical treatment of hip instability in patients with lower lumbar level myelomeningocele: is muscle transfer required? *Clin Orthop Relat Res.* 2015; 473(10):3254-60.
50. Menelaus MB. Progress in the management of the paralytic hip in myelomeningocele. *Orthop Clin North Am.* 1980; 11(1):17-30.
51. Buckley SL, Sponseller PD, Magid D. The acetabulum in congenital and neuromuscular hip instability. *J Pediatr Orthop.* 1991; 11(4):498-501.
52. Mannor DA, Weinstein SL, Dietz FR. Long-term follow-up of Chiari pelvic osteotomy in myelomeningocele. *J Pediatr Orthop.* 1996; 16(6):769-73.
53. Zenios M, Hannan M, Zafar S, Henry A, Galasko CS, Khan T. Clinical and radiological outcome of combined femoral and Chiari osteotomies for subluxed or dislocated hips secondary to neuromuscular conditions: a minimum of 10-year follow-up. *Musculoskelet Surg.* 2012; 96(2):101-6.
54. Breed AL, Healy PM. The midlumbar myelomeningocele hip: mechanism of dislocation and treatment. *J Pediatr Orthop.* 1982; 2(1):15-24.
55. Abel MF, Sutherland DH, Wenger DR, Mubarak SJ. Evaluation of CT scans and 3-D reformatted images for quantitative assessment of the hip. *J Pediatr Orthop.* 1994; 14(1):48-53.