

PERSPECTIVE

Iranian Joint Registry

(Iranian National Hip and Knee Arthroplasty Registry)

Hamidreza Aslani, MD; Seyed Taghi Nourbakhsh, MD; Farivar A. Lahiji, MD; Keykavoos Heydarian, MD; Mahmood Jabalameli, MD; Mohammad Taghi Ghazavi, MD; Mohammad Naghi Tahmasebi, MD; Mahmoud Reza Fayyaz, MD; Mohammad Ali Sazegari, MD; Maziar Mohaddes, MD; Mojtaba Rajabpour, MD; Mohammad Emami, MD; Seyyed Mohammad Jazayeri, MD; Firooz Madadi, MD; Hossein Farahini, MD; Fardin Mirzatoolee, MD; Mohammad Gharahdaghi, MD; Mohammad Hossein Ebrahimzadeh, MD; Mohammadreza Ebrahimian, MD; Hossein Mirvakili, MD; Kaveh Bashti, MD; Mohtasham Almasizadeh, MD; Mansour Abolghasemian, MD; Afshin Taheriazam, MD; Mehdi Motifard, MD; Hamidreza Yazdi, MD; Mahmood Karimi Mobarakeh, MD; Masoud Shayestehazar, MD; Mehdi Moghtadae, MD; Babak Siavashi, MD; Mohammadreza M. Sajjadi, MD; Alireza Manafi Rasi, MD; Seyyed Kazem Chabok, MD; Zohreh Zafarani, MD; Shahin Salehi, MD; Monireh Ahmadi, PhD; Amin Mohammadi, MD; Mohammad Ebrahim Shahsavand, MD

Research performed at Shahid Beheshti University of Medical Sciences, Knee and Sport Medicine Education and Research Center, Tehran, Iran

Received: 3 February 2016

Accepted: 25 March 2016

Abstract

Periodic evaluation and monitoring the health and economic outcome of joint replacement surgery is a common and popular process under the territory of joint registries in many countries. In this article we introduce the methodology used for the foundation of the National Iranian Joint Registry (IJR) with a joint collaboration of the Social Security Organization (SSO) and academic research departments considering the requirements of the Iran's Ministry of Health and Education.

Keywords: Arthroplasty, Hip, Iran, Knee, Registry national

Introduction

Total joint arthroplasty is a common procedure and its safety and effectiveness has been proved in many papers (1-4). Between 1990 and 2002 the rate of total hip arthroplasty increased by 50% and the rate of primary total knee arthroplasty almost tripled (5, 6). Exponential rise in primary arthroplasties is expected to double the rate of revision surgeries in the next two decades (6). Anticipated rise in arthroplasty requires a reliable and valid method of evaluation and monitoring the safety and effectiveness in the region (7-9).

Registries for hip and knee replacements are becoming popular worldwide with growing importance. Many of the preeminent registries – such as those in Sweden, Finland, Norway, Australia, Denmark, and New Zealand – have more than 10 years of experience and are currently collecting nationwide data on more than 90 percent of procedures (7, 10-13).

Identifying the epidemiology of the joint arthroplasty, evaluating the outcome of the surgery and collecting timely information of the risk factors associated with poor outcome are the main goals of the registries (14).

Moreover, registries had an important role in improving our knowledge (9, 14). It has been shown a reduction in the number of revision hip replacement surgeries in Australia from 13 percent in 2003 to 11.2 percent in 2010 (15). In the U.S., revision rate was 16.9 for patients underwent primary total hip replacement from 1992 to 2002 although they lack an integrated registry (5).

Reducing the rate of revision is not only a sign of better clinical outcome but also dramatically influences the economic burden of the disease on patients and health care systems. Each percentage point reduction in revision surgeries saves an estimate of \$42.5 million to \$112.6 million annually (According to the national estimates of the United States) (5).

We were not able to find any report about the rate of arthroplasty in Iran. National Social Security organization (SSO) as one of the most important with extensive health insurance coverage in Iran decided to jointly develop a national joint arthroplasty registry. Reduction in complications while improving the outcomes after joint arthroplasties, in addition to reducing the economic burden of this procedure are the

Corresponding Author: Hamidreza Aslani, Shahid Beheshti University of Medical Sciences, Knee and Sport Medicine Education and Research Center, Tehran, Iran
Email: hraslani@yahoo.com



THE ONLINE VERSION OF THIS ARTICLE
ABJS.MUMS.AC.IR

main goals of this registry. In this study, we introduce our methodology in the development of the National Iranian Joint Registry (IJR) with a joint collaboration of the Social Security Organization (SSO), considering the requirements of this organization as well as the Iran's Ministry of Health and Education.

Material and Methods

Knee and Sports Medicine Research Center (KSMRC) began to develop a national joint arthroplasty registry in June 2014 according to the mission requested by the SSO. Moreover, this center is responsible for the establishment and implementation of this system in a population under the coverage of SSO insurance, which then will be extended to the whole nation.

We used four sources of information for the design and development of seven stages of the Iranian Joint Registry (IJR) [Figure 1].

1. Interview with the policy makers of the SSO, other insurance providers, and the ministry of health providers: We met with six member of the Supreme Council of the SSO. All conversations were recorded and documented.

2. Review of the reports from international registries: We included eight registries in our review including registries of Sweden, Finland, Norway, England, Italy, France, Germany, and Hungary.

3. Site visit: Visiting the Swedish Hip Arthroplasty Registry site

4. Experts meetings: Six experienced orthopedic surgeons, official members of the IOA (Iranian Orthopedic association), determined the scientific and indigenous issues in regards to the IJR over ten meetings with a focused group discussion method.

Data and information from these resources was used to



Figure 1. Iranian Joint Registry (IJR).

develop the IJR process through seven stages:

1. Developing the Minimal Data Set
2. Development of the process for data collection
3. Development of the process for collecting the registered data and timely feed back
4. Development of the leadership, management and funding procedure.
5. Legislation and Protocols
6. Development of strategies for evaluation and validation of the IJR
7. Development of strategies for expanding the registry to a national level

Result

IJR was developed in seven stages

1. Minimal Data set

Patients, surgeon providers, patients' insurance status, anesthesia, indications, preoperative deformities, core surgical date, thromboprophylaxis regimens, implant data, application of the bone graft, and intra operative events are the essential primary data, which are included in the IJR forms. We developed six forms, three for the knee and three for the hip joint [Figure 2]. There is one form for primary replacement surgeries (K1 and H1), there are two forms for the second stage or the revision surgeries (stage 1, 2), and one form for one stage revision surgeries.

1. Data collection Procedure

Data entry is electronic. Patients' information will be entered according to the hospital information system (HIS) by the operation room assistant (step 2). Other details of surgery and treatment will be entered by the surgeon provider [Figure 3]. The software is designed to transfer the data to the operation note at the completion of data entry.

1. Collecting the registered data and timely feed back

Online data entry in the IJR helps receive a timely feedback. Knee and Sports Medicine Research Center




Figure 2. Data entry fom classification in IJR.

(KSMRC) is responsible for the annual analysis and report of the data. Reporting the survival rate of the implants and factors associated with early revision are the primary aims of the IJR.

2. Leadership, Management and Funding procedure

SSO is providing the major funding support to the IJR while KSMRC is responsible for implementation and maintenance of the IJR and collaboration with the other hospital.

3. Legislation and Protocols

Currently completing the IJR forms are arbitrary for surgeons. However, there is a planned legislation, which will obligate surgeons to register the procedure in order to be reimbursed.

4. Evaluation and validation strategy

SSO insurance evaluates and monitors the monetary costs to reimburse the hospitals and providers. IJR forms will be under the evaluation list of the SSO as part of the paperwork required for being reimbursed. KSMRC will analyze the registered data every 3 month. Surgeons will be asked to complete the data entry if discrepancies or missed data are found.

5. Strategies for expanding the registry to a national level

In the first step, Milad Hospital in Tehran - affiliated with the SSO - will launch the IJR as an obligatory assignment for the providers. In parallel, several university hospitals will start registration under voluntary circumstances. In the next step, completing the forms for patients who

are under the coverage of the SSO insurance will become mandatory. In other steps, we will collaborate with other university based hospitals and health insurance organizations to expand the coverage of the IJR.

Discussion

Arthroplasty registries are employed in several countries where the growing prevalence of replacement surgeries highlights the importance of registries. Development of the Iranian joint registry helps reduce the morbidity as well as reducing the cost of the procedure for patients. In this article we introduced the design of the arthroplasty registry following seven stages. We hope to reach to a valid and reliable national registry in a timely manner.

The IJR's minimal data set is created according to recommendations from the International Society of Arthroplasty Registries (ISAR) (16). Data submission and data collection differs between registries (7). In Finland, Italy and Norway the data is submitted using paper forms. Australian and Canadian registries use both paper and electronic forms (7,15). The IJR will be based on a single digital platform. The software will transfer the entered data to the operation note automatically. Adding this step will make the data entry less cumbersome for the providers.

Governmental organizations are commonly found to be the initiator of the registries in other countries. However,




Figure 3. Data entry processes in IJR.

to our best knowledge it is the first time that non-governmental insurance organization has been involved in initiating a national joint replacement registry.

Iranian joint registry is designed, which will be developed to form the major source of outcome assessment after orthopedic surgeries. Joint collaboration of the national organizations is an essential requirement for the advancement of this mission.

Acknowledgment

Iran national social security organization and different Iranian universities supported this research. We thank our colleagues from Iranian Orthopedic Association, who provided insight and expertise that greatly assisted the research, although they may not agree with all of the interpretations/conclusions of this paper. We would like to thank all Iranian universities orthopedic department for their help and support. We do appreciate for help of Management Committee Members and orthopedic department of Shahid Beheshti university of medical science for their encouragement and their support. We do appreciate Swedish hip registry especially Professor Göran Garellick, Register Director Maziar Mohaddes, University of Gothenburg Department of Orthopaedics. Consultant in Orthopaedics at Sahlgrenska University Hospital & Kajsa Erikson Register Coordinator for their hospitality and friendly communication, support, and help.

Also we regarded special thanks to Milad Hospital Board of Directors, Orthopedic Department and technical supports of Computer and IT group.

Hamidreza Aslani MD
Shahid Beheshti University of Medical Sciences, Knee and Sport Medicine Education and Research Center, Tehran, Iran

Seyed taghi Nourbakhsh MD
Mahmoud Reza Fayyaz MD
Mohammad Ali Sazegari MD
Department of Orthopedic Surgery, Milad Hospital, Knee and Sport Medicine Education and Research Center, Tehran, Iran

Farivar A. Lahiji MD
Mohammad Emami MD
Seyyed Mohammad Jazayeri MD
Firooz Madadi MD
Mohammadreza M. Sajjadi MD
Alireza Manafi Rasi MD
Department of Orthopedic Surgery, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Keykavoos Heydarian MD
Mahmood Jabalameli MD
Mohammad Taghi Ghazavi MD
Hossein Farahini MD
Mansour Abolghasemian MD
Hamidreza Yazdi MD
Mehdi Moghtadae MD
Department of Orthopedic Surgery, Iran University of Medical Sciences, Tehran, Iran

Mohammad Naghi Tahmasebi MD
Kaveh Bashti MD
Babak Siavashi MD
Department of Orthopedic Surgery, Tehran University Medical Sciences, Tehran, Iran

Maziar Mohaddes MD, PhD
Department of Orthopaedics, Institute of Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Sahlgrenska University Hospital

Mojtaba Rajabpour MD
Clinical Research Center Milad Hospital, Knee and Sport Medicine Education and Research Center, Tehran, Iran

Fardin Mirzatoloe MD
Department of Orthopedic Surgery, Urmia University of Medical Sciences, Urmia, Iran

Mohammad Gharahdaghi MD
Mohammad Hossein Ebrahimzadeh MD
Orthopedic Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

Mohammadreza Ebrahimian MD
Hossein Mirvakili MD
Mohtasham Almasizadeh MD
Department of Orthopedic Surgery, Milad Hospital, Tehran, Iran

Afshin Taheriazam MD
Department of Orthopedic Surgery, Tehran Medical Sciences Branch, Islamic Azad University, Tehran, Iran

Mehdi Motifard MD
Department of Orthopedic Surgery, Isfahan University of Medical Sciences, Isfahan, Iran

Mahmood Karimi Mobarakeh MD
Department of Orthopedic Surgery, Kerman University of Medical Sciences, Kerman, Iran

Masoud Shayestehazar MD
Orthopedic Research Center, Mazandaran University of Medical Sciences, Sari, Iran

Seyyed Kazem Chabok MD
Consultant Orthopedic Surgeon, Ordibehesht Hospital, Shiraz, Iran

Shahin Salehi MD
Department of Sports Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Zohreh Zafarani MD
Monireh Ahmadi PhD
Amin Mohamadi MD, MPH
Mohammad Ebrahim Shahsavand MD
knee and Sport Medicine Research and Education Center Milad Hospital, Tehran, Iran

References

1. Callaghan JJ, Beckert MW, Hennessy DW, Goetz DD, Kelley SS. Durability of a cruciate-retaining TKA with modular tibial trays at 20 years. *Clin Orthop Relat Res.* 2013; 471(1):109-17.
2. Santini AJ, Raut V. Ten-year survival analysis of the PFC total knee arthroplasty--a surgeon's first 99 replacements. *Int Orthop.* 2008; 32(4):459-65.
3. Kachooei AR, Claessen FM, Chase SM, Verheij KK, van Dijk CN, Ring D. Factors associated with removal of a radial head prosthesis placed for acute trauma. *Injury.* 2016; 1383(16):30022-5.
4. Kachooei AR, Chase SM, Jupiter JB. Outcome assessment after aptis distal radioulnar joint (DRUJ) implant arthroplasty. *Arch Bone Jt Surg.* 2014; 2(3):180-4.
5. Kurtz S, Mowat F, Ong K, Chan N, Lau E, Halpern M. Prevalence of primary and revision total hip and knee arthroplasty in the United States from 1990 through 2002. *J Bone Joint Surg Am.* 2005; 87(7):1487-97.
6. Kurtz SM, Lau E, Ong K, Zhao K, Kelly M, Bozic KJ. Future young patient demand for primary and revision joint replacement: national projections from 2010 to 2030. *Clin Orthop Relat Res.* 2009; 467(10):2606-12.
7. Serra-Sutton V, Allepuz A, Espallargues M, Labek G, Pons JM. Arthroplasty registers: a review of international experiences. *Int J Technol Assess Health Care.* 2009; 25(1):63-72.
8. Bourne RB. The planning and implementation of the Canadian Joint Replacement Registry. *Bull Hosp Jt Dis.* 1999; 58(3):128-32.
9. Maloney WJ. National Joint Replacement Registries: has the time come? *J Bone Joint Surg Am.* 2001; 83-A(10):1582-5.
10. Karrholm J. The Swedish Hip Arthroplasty Register (www.shpr.se). *Acta Orthop.* 2010; 81(1):3-4.
11. Makela KT, Matilainen M, Pulkkinen P, Fenstad AM, Havelin LI, Engesaeter L, et al. Countrywise results of total hip replacement. An analysis of 438,733 hips based on the Nordic Arthroplasty Register Association database. *Acta Orthop.* 2014; 85(2):107-16.
12. Havelin LI. The Norwegian Joint Registry. *Bull Hosp Jt Dis.* 1999; 58(3):139-47.
13. Lucht U. The Danish Hip Arthroplasty Register. *Acta Orthop Scand.* 2000; 71(5):433-9.
14. Gioe TJ, Killeen KK, Mehle S, Grimm K. Implementation and application of a community total joint registry: a twelve-year history. *J Bone Joint Surg Am.* 2006; 88(6):1399-404.
15. Owen DH, Russell NC, Smith PN, Walter WL. An estimation of the incidence of squeaking and revision surgery for squeaking in ceramic-on-ceramic total hip replacement: a meta-analysis and report from the Australian Orthopaedic Association National Joint Registry. *Bone Joint J.* 2014; 96-B(2):181-7.
16. Erikson K. 5th International Congress of Arthroplasty Registries, Wrightington and Manchester, UK. ISAR, International Society of Artbroplasty Registers. available at: URL: <http://www.isarhome.org/>; 2016.